

Simulating the collaborative cross: power of QTL detection and mapping resolution in large sets of recombinant inbred strains of mice

William Valdar, Jonathan Flint, Richard Mott

Abstract

It has been suggested that the collaborative cross, a large set of recombinant inbred strains derived from eight inbred mouse strains, would be a powerful resource for the dissection of complex phenotypes. Here we use simulation to investigate the power of the collaborative cross to detect and map small genetic effects. We show that for a fixed population of 1,000 individuals, 500 RI lines bred using a modified version of the collaborative cross design are adequate to map a single additive locus that accounts for 5% of the phenotypic variation to within 0.96 cM. In the presence of strong epistasis more strains can improve detection, but 500 lines still provide sufficient resolution to meet most goals of the collaborative cross. However, even with a very large panel of RILs, mapping resolution may not be sufficient to identify single genes unambiguously. Our results are generally applicable to the design of RIL in other species.

Introduction

The Complex Trait Consortium (CTC) outlined a strategy, called the Collaborative Cross (CC), to construct a very large set of recombinant inbred (RI) strains from a genetically diverse set of inbred strains of mice (CHURCHILL *et al.* 2004). It is suggested that the CC is an ideal resource for systems biology as it will provide a reproducible, highly varied yet controlled set of genetic backgrounds for functional genomic studies. One projected use of

the CC is the detection and mapping of genetic effects underlying complex phenotypes. In principal, the CC could solve the obdurate problem of high resolution quantitative trait locus (QTL) mapping: the CTC position paper suggested the creation of 1,000 RIs to map QTLs to a resolution where gene identification becomes possible, within a reasonable time and budget. An international collaboration to breed these lines has started, and is described at <http://www.complextait.org>.

However, the experimental design and statistical power of the CC has not yet been rigorously assessed. There are a number of factors that need to be considered. First, the genetic architecture of complex traits, which remains largely unknown, will influence the CC's ability to detect and resolve genetic effects. Second, it is not clear how best to design the cross to maximize and maintain genetic diversity. If the CC is not seeded with sufficient diversity in its initial stages, or loses diversity during breeding, there may be insufficient detectable recombinants in the RI strains to deliver adequate mapping resolution. Third, assuming that the CC does have the potential to deliver high resolution QTL mapping, how does it compare to other possibly more efficient strategies? For instance, detection and fine-mapping can be achieved within a population of manageable size by making use of historical recombinants. Advanced intercross lines (AIL), populations made from F₂s by repeated intercrossing within a large population, have both the simple genetic background necessary for unambiguous assignment of identity by descent along with the highly recombinant

haplotype structure that allows high mapping resolution (DARVASI and SOLLER 1995; WANG *et al.* 2003). Heterogeneous stocks (HS) are a generalization of AIL, derived from a cross of eight inbred mouse lines (DEMAREST *et al.* 2001; MCCLEARN *et al.* 1970). Although HS are more complex to analyse, they provide additional genetic diversity and may offer increased mapping resolution over AILs (TALBOT *et al.* 2003). But AIL and HS have drawbacks, since they take years to breed and are expensive to genotype. The efficiency of AIL and HS designs, relative to the CC, has not been investigated.

In this paper, we use simulation to compare QTL detection and accuracy of the collaborative cross design against several alternatives. We simulate the realistic situation of a quantitative trait that owes half its variance to genetic factors. Of that half, a small portion is due to a QTL on a genotyped chromosome. The remaining variance is attributable to multiple QTLs that lie on ungenotyped, hence unobserved, chromosomes. We investigate the ability of the CC to detect and map a QTL when it acts independently and when it is under strong epistatic control. We investigate variations on the basic breeding design, compare it with traditional alternatives (F2 intercross and the backcross) and newer, related strategies (RIHS (VALDAR *et al.* 2003) and RIs based on AILs). Finally we ask how many lines are necessary to fine-map small effect QTLs both in simple additive and in strongly epistatic systems.

Methods

Breeding strategies overview

We examined five breeding strategies: the Collaborative Cross (CC) ((THE COMPLEX TRAIT CONSORTIUM), (CHURCHILL *et al.* 2004)), recombinant inbred heterogeneous stocks (RIHS) (VALDAR *et al.* 2003), recombinant inbred advanced intercross lines (RIAIL), the F2 intercross (F2) and the backcross (BC). RIHS and RIAIL are panels of recombinant inbred strains made from respectively HS and AI populations. With the exception of the backcross, these strategies are variants of a more general multistage breeding programme. This programme has three stages: a compulsory mixing stage, followed by two optional stages, maintenance and inbreeding (Figure 1). In the mixing stage, inbred progenitor strains are intercrossed to produce a foundation population of mice whose genomes each contain some genetic material from every founder strain. In the maintenance stage, foundation mice are intercrossed over a number of generations. The aim is to introduce recombinants and produce a population of mice whose genomes are fine-grained mosaics of the original founder haplotypes. Intercrossing in the maintenance stage is typically performed using a pseudo-random mating technique that avoids pairing of close relatives and thereby reduces genetic drift and the risk of fixation. In the inbreeding stage, randomly chosen pairs of mice are inbred by recurrent brother-sister mating over 20 generations, each pair producing a distinct recombinant inbred line. We describe the four main types of strategies, and their variants, below and in Table 1.

F2 intercross and backcross

In an F2 cross two inbred lines, A and B, are crossed in the mixing stage, to make F1 foundation animals, of genetic composition AB. They enter a maintenance stage of one generation to produce F2 animals. The inbreeding stage is omitted. For the backcross, two inbred lines, A and B, are combined in an F1 cross. These foundation animals are then crossed with strain A to produce animals whose chromosome pair comprises one A haplotype and one recombinant AB haplotype.

Recombinant inbred advanced intercross strains (RIAILs)

Advanced intercross lines (DARVASI and SOLLER 1995), resemble an F2 with an extended maintenance stage that lasts two or more generations. AIL genomes thus also have genetic composition AB, but contain more recombinants. To provide an informative comparison with the CC and RIHS crosses, which is the true focus of our work, we do not simulate AILs as such, but instead recombinant inbred strains derived from an AIL (RIAIL), made by adding on an inbreeding stage to the traditional AIL design. The genomes of RIAIL mice are therefore highly recombinant mosaics of strains A and B that theoretically are homozygous and identical within a particular line but different between lines. We use a notation in which RIAILs are suffixed with the number of filial generations intercrossed before inbreeding, eg, RIAIL25 means RIs produced from an F25 or, equivalently, an AIL25 population.

Recombinant inbred heterogeneous stocks (RIHS)

Recombinant inbred heterogeneous stocks (VALDAR *et al.* 2003) are like RIAIL but derived from eight founders rather than two. They are made as follows. Eight inbred lines, A to H, are combined in a three-step mixing phase. In the first step, four F1 populations are produced of compositions AB, CD, EF and GH by mating A with B, C with D and so on, i.e. a “two-way” cross. The second step proceeds with two “four-way” crosses: AB and CD mice are mated to produce mice of composition ABCD, while EF and GH similarly give rise to EFGH. The last step is an “eight-way” cross, in which ABCD mice are crossed with EFGH mice to produce mice of genetic composition ABCDEFGH. We call these animals foundation heterogeneous stocks, or HS0. After maintenance for n generations to build up recombinants they become HS n . Subsequent inbreeding produces RIHS n , e.g. RIHS30 are RIs produced by inbreeding pairs of HS30 mice.

Collaborative Cross (CC)

The Collaborative Cross as initially proposed (CHURCHILL *et al.* 2004; THE COMPLEX TRAIT CONSORTIUM) is much like RIHS but with a more elaborate mixing stage and no maintenance breeding. In its most extreme form, mixing is ambitiously complex. Simpler mixings have since been suggested (CHURCHILL *et al.* 2004), whose complexity lies between that of the original and that of simple RIHS mixing. Here we simulate the most extreme form, noting that this may overestimate the efficacy of an intermediate. As in RIHS, the mixing stage of the Collaborative Cross has three steps. The first step produces F1s from every distinct pair

of founders, generating mice of composition AB, AC, ..., FH, GH. This “diallel” design comprises 28 two-way crosses, or 56 if reciprocal crosses between males and females are performed (which they are not here as we restrict our attention to autosomes). In step two, every possible four-way cross is performed between F1s excluding instances of shared descent. For example, AB x CD is allowed but AB x BC is not. We call this a “disjoint diallel”. In the last step, eight-way crosses are performed in a further disjoint diallel (where, for example, ABCD x EFGH is allowed but ABCD x ABDE is not) resulting in foundation mice of composition ABCDEFGH. Its repeated disjoint diallel structure means the mixing stage incurs a combinatorial explosion of crosses. In the unrealistically conservative situation of every distinct cross being performed with no more than one mating pair, step two would see 210 matings and step three, 310. Only a subset of those eight-way crosses would be feasible in practice. We simulate them all.

In the standard CC proposal, pairs of foundation mice then are inbred to produce RIs. We also investigated variant CC designs with more recombination, in order to increase mapping accuracy. In “cctop10”, we select only the most highly recombinant 10% of mice from the four-way cross (which could be determined by genotyping) to proceed to the eight-way cross. In “cc04” and “cc08”, we insert four and eight maintenance generations (about one and two years of extra breeding) respectively between the mixing and inbreeding.

Maintenance stage

There are several established ways to maintain a laboratory population over many generations with a view to preserving heterozygosity (CABALLERO and TORO 2000). Circular mating (also known as rotational breeding) defines a structured mixing of the animals, repeated at each generation that implicitly restricts the consanguinity of each mating and so minimizes the extent of inbreeding, drift and fixation (BOUCHER and COTTERMAN 1990; KIMURA and CROW 1963). We perform maintenance in all strategies using circular mating. We simulate it as follows. Given a population of $2m$ individuals, split equally between the sexes, put couples into cages labelled 1 to m . Prepare empty cages labelled 1' to m' to hold the next generation. Mate the female from cage 1 with the male from cage 2 and send their offspring to cage 1'. In similar fashion, mate the female of cage k with the male from cage $k+1$ (or the male from cage 1 if $k=m$) and send their offspring to cage k' , doing this for all remaining k . Repeat this procedure, with suitable relabelling, until the required number of maintenance generations is reached.

The genetic diversity of the final population depends critically on m , the number of mating pairs. When m is small, the maintenance stage acts as a population bottleneck that encourages extremes of drift and fixation. Conversely, a larger m delays these effects. Consistent with existing HS mice (Northport HS, (DEMAREST *et al.* 2001)), we set m at 50 when generating RIHS lines unless otherwise specified, and do this for RIAIL also. However, for our CC

variants, project of more ambitious scope, we set m equal to the number of RIs required, which can be as high as 1,000.

Founder strains and marker sets

Northport HS stocks derive from a cross of the mouse strains A/J, AKR/J, BALB/cByJ, C3H/HeJ, C57BL/6J, CBA/J, DBA/2J and LP/J. Hitzemann chose those strains with dual purpose: to emulate the original Boulder HS, whose founders were genetically diverse by the standards of that day, and to provide a contrast of response classes to haloperidol-induced catalepsy (Robert Hitzemann, personal communication).

The Complex Trait Consortium urge that the founder set chosen for the CC balances genetic diversity, phenotypic diversity, breeding performance and the availability of complementary resources (THE COMPLEX TRAIT CONSORTIUM). The strains finally selected for breeding by the CTC were A/J, C57BL/6J, 129Sv/ImJ, NOD/LtJ, NZO/H1J, CAST/Ei, PWK/Ph, WSB/Ei, which includes a number of wild-derived strains. However, at the time of simulation this set was not yet finalized, and only incomplete genotype data was available for them. For these reasons, and because we wished to compare alternative eight-founder designs on an equal footing, all simulated populations were founded with the Northport set listed above, or, in the case of the two-strain strategies, with pairs of strains drawn from that list. The genomes of wild-derived strains such as CAST/Ei are more distantly related than are standard strains such as C57BL/6J. Therefore we would expect more QTLs to segregate in the actual CC but

the results of our simulations should not change overmuch since they are conditioned on the presence of segregating QTLs.

We use two marker sets. The first is an artificial fine-grain set of 1,000 fully informative markers (ie, markers with eight alleles) spaced at regular intervals of 0.1 cM. We use this set only to establish unrecombined block length for different designs (see later) and not for mapping. The second, which we use to map QTLs, is a real dataset typed for chromosome one, provided by the Genomics Institute of the Novartis Research Foundation (GNF) (PLETCHER *et al.*). We selected markers that segregate among the eight Northport HS progenitors in order to make our simulations as realistic as possible. The GNF set numbers 512 diallelic SNPs with average spacing 0.19 cM, (standard deviation 0.16 cM) over 100 cM. These markers segregate among the eight founders with varying minor allele frequencies. The strain distribution pattern of the SNPs are such that of the 512 minor alleles, 68 are present in only one strain, 136 are present in two, 209 in three and 99 in four.

Representing the mouse genome

We represent a mouse by up to seven pairs of homologous autosomes, depending on the complexity of the genetic model being simulated . Chromosome 1 is 100 cM long and contains the GNF marker set, or in the case of establishing block lengths the fine-grain artificial marker set. The QTL to be mapped always lies on Chromosome 1. The remaining six chromosomes act as an unseen genetic background that affects the phenotype; these

chromosomes are ignored during mapping. For our simulations of epistasis, Chromosome 2 contains a single diallelic marker that acts as a masking QTL in our simulation of epistasis but is ignored otherwise. Chromosomes 3-7 each contain 10 diallelic markers spaced at 10 cM intervals. These 50 markers are used as background QTLs (see below). Each of the 51 markers on chromosomes 2-7 has its high allele on four founders independently and randomly chosen from the eight.

Simulating breeding in the mouse population

We simulate mouse populations stochastically using software written in-house (available from <http://www.well.ox.ac.uk/~valdar/software/>). Gametogenesis is simulated by randomly drawing one of each parental autosome and recombining it with its homolog, choosing breakpoints according to the Haldane model with no interference, in which meioses occur as events in a one dimensional Poisson process (LYNCH and WALSH 1998).

Phenotypic variance components

We simulate a quantitative phenotype that is 50% heritable in the founder population. This means genetic factors explain half the phenotypic variance and environmental noise accounts for the remainder. The heritable portion of the variance is split between foreground and background QTLs. Foreground QTLs are defined as those that lie on chromosome 1. These QTLs are visible in our mapping. Background QTLs lie on the unseen chromosomes (2-7)

and, although they affect the phenotype, are unmappable. The phenotypic variance partitions thus:

$$\begin{aligned}\sigma_P^2 &= \sigma_G^2 + \sigma_E^2 \\ &= (\sigma_Q^2 + \sigma_B^2) + \sigma_E^2\end{aligned}$$

where subscripts identify phenotypic (P), environmental (E), genetic (G), foreground (Q) and background (B) variance components, and where, for a phenotype that is 50% heritable,

$$\sigma_G^2 = \frac{1}{2}\sigma_P^2.$$

QTL effect sizes and allelic values

The effect size of a QTL is defined here as the proportion of phenotypic variance it explains in the founder population, eg, $\theta_Q = \sigma_G^2 / \sigma_P^2$. We explicitly vary the effect size of a single foreground QTL while implicitly adjusting the effect sizes of the background QTLs to make up the remaining heritability. These 50 background QTLs act independently to provide only additive genetic variation. Thus, ignoring covariance between QTLs (which is negligible in our design), $0.5 = \theta_Q + \sum_{i=1}^{50} \theta_{B_i}$, where the effect size of the i th background QTL is $\theta_{B_i} = (0.5 - \theta_Q) / 50$, such that in any one simulation all background QTLs are equally potent in the founder population. The case of epistasis is described below.

The allelic value is defined here as the raw number that a QTL's high allele contributes to the phenotype value. We adjust each QTL's allelic value at the start of each simulation to yield the required effect size in the founder population. That allelic value then remains constant throughout breeding and phenotyping. This means, owing to inevitable drifts in allele frequency, a QTL's effect size in the mapping population may differ from its starting value in the founder population, but it models reality better than fixing the effect size in the mapping population by changing its allelic value.

Simulating QTLs

We simulate two main foreground QTL systems: a single additive QTL, and a single QTL in strong epistasis with an unseen QTL on chromosome 2. In both cases, background QTLs are present but act independently of QTLs in the foreground.

We simulate a single foreground QTL as follows. Before any breeding takes place, a marker is drawn at random from chromosome 1 and designated the QTL. Then the strain distribution pattern (SDP) of its alleles is shuffled. We do this for the following reason. In the GNF marker set, 90% of the SNPs segregate between C57BL/6 and A/J or DBA/2, due to the method of SNP ascertainment. As a result, the strain distribution pattern of the QTL matches that of nearby SNPs more frequently than would be likely in reality, since there is no reason to suppose a QTL favours any particular SDP. To correct for this bias, we shuffle the SDP of the QTL but keep the SDP of the remaining SNPs unchanged. For example, a QTL with its

minor allele in strains A, B and C may, after shuffling, find that allele moved to strains B, F and H. Were the strain distribution pattern of the QTL not shuffled it would be easy to detect through single marker association because its allelic state would be closely matched by those of its flanking markers. We demonstrate this point by performing an additional set of unshuffled simulations for one experimental condition, contrasting its results with those of the shuffled case (see Results). Once phenotyping is complete, the foreground QTL marker is concealed, and we then attempt to recover that marker's location.

The high allele of each QTL is assigned an allelic value based on its frequency and required effect size in the founder population. The phenotype of each animal in the mapping

population is calculated as $y = \alpha_Q x_Q + \sum_{B=1}^{50} \alpha_{B_i} x_{B_i} + \varepsilon$, where, with subscripts B identifying

background loci, α_B denotes the allelic value, x_B the number of high alleles, and ε a normally distributed error term with variance accounting for half the total variance in the founders. Table 2 illustrates the relation between the genotype (x) of an additive locus and its genetic effect (αx). For inbred mice x is usually 0 or 2 (though can be 1 if inbreeding has not fixed that QTL), for F2 mice x is 0, 1 or 2, and for BC mice x is 0 or 1.

For simulations of epistasis, the foreground and background QTLs are generated as before except that the genetic affect of the foreground QTL on chromosome 1 now depends on the masking QTL, M , on chromosome 2. The equation for a mouse phenotype is now

$y = \alpha_{Q \times M} x_M x_Q + \sum_{i=1}^{50} \alpha_{B_i} x_{B_i} + \varepsilon$, such that, for example, in a population of inbred lines the genetic effect from the foreground QTL ($\alpha_{Q \times M} x_M x_Q$) is non-zero only when both foreground and masking QTLs are in the high state. Note that the “effect size” of the foreground QTL is now defined as the proportion of variance explained by the QTL pair. The goal of subsequent detection and fine mapping is to find the first QTL despite the other’s confounding effect.

The foreground and masking QTLs always segregate in each simulation. Thus for F2, RIAIL and BC, founded on just two strains, the pair of strains chosen is random, subject to the constraint that those QTLs segregate between them.

Detection and fine-mapping

We detect and fine-map simulated QTLs using two methods, single marker association (SMA) and HAPPY, a multipoint method (MOTT *et al.* 2000). In single marker association, we regress the phenotype on the number of minor alleles present at each marker in turn. If the marker showing the strongest association is statistically significant (see later), we classify the QTL as detected and predicted to be located in the most significant marker interval. Otherwise, we deem the QTL undetected.

Mapping using HAPPY is also performed in a regression framework, but here the phenotype is regressed on the inferred probability of descent at each marker interval. Briefly, HAPPY models genotypes as symbols emitted from hidden progenitor states in a discrete Markov process where transitions between states correspond to recombination events. This hidden Markov model (HMM) is fitted to the data using a dynamic programming algorithm described in (MOTT *et al.* 2000). The HMM transition matrix is parameterized by the expected number of recombinants between markers, which, given the centiMorgan distance between markers, can be summarized by the effective number of generations g_e . Inbreeding, among other things, causes g_e to deviate from the actual number of generations and so g_e varies with breeding strategy. For each strategy we estimate g_e by calibrating the observed level of informative recombination against that seen in a simulated outbred population.

By combining all the genotype data along a diploid chromosome, HAPPY computes for each individual and at locus the probability of descent from each pair of the founder strains. The progenitor probabilities are regressed on the phenotype to identify loci where the pattern of strain inheritance correlates with the phenotype. As with SMA, the marker interval that associates most strongly with the phenotype is tested for statistical significance and, if it passes, declared the putative QTL.

In each simulation, 1,000 animals are generated. As a consequence, in some experiments multiple animals are phenotyped from each RI strain. In that case we regress the mean phenotype of those replicates (ie, the “strain mean”) on genetic information from a randomly chosen member. Residual heterozygosity within an RI means “replicates” are rarely identical at all loci, so we implicitly model the drawbacks of minimal genotyping.

Assessing statistical significance using Generalized Extreme Value distributions

We measure the association between locus and phenotype in both single marker association and HAPPY by an F-test of their respective linear model fits. The overall score for a simulation over m loci is defined as follows. Let $S_i = -\log_{10} P_i$, where P_i is the p-value from the F-test at locus i . Define $S_{\max} = \max\{S_1, \dots, S_m\}$, that is, the maximum score over all loci. The reported p-value for the highest scoring locus is grossly anticonservative. We therefore assess statistical significance by comparing the obtained p-value with genomewide thresholds that have been corrected for multiple testing. The Bonferroni correction, is too conservative as it ignores linkage disequilibrium (LD) between markers, and a permutation test is computationally unfeasible for so many simulations. Instead, we take LD into account by modelling the distribution of S_{\max} under the null hypothesis of no foreground QTL and hence calculate suitable threshold quantiles.

The LD structure varies with breeding strategy, number of RI lines and extent of background genetic variation so we estimate the null distribution of S_{\max} separately for each experimental

condition. Provided there is negligible long-range dependence between distant S_i , with increasing m , S_{\max} asymptotically follows a Generalized Extreme Value distribution (COLES 2001) with distribution function $G(S_{\max}) = \exp\left\{-\left[1 + \xi(S_{\max} - \mu)/\sigma\right]_+^{-1/\xi}\right\}$, where μ , σ and ξ are the location, scale and shape parameters respectively, and $h_+ = \max(h, 0)$. A closely related use of GEVs is described in ref (DUDBRIDGE and KOELEMAN 2004).

For each experimental condition we perform 1,000 null simulations with background QTLs only. We then fit a GEV by maximum likelihood to the observed values of S_{\max} using the R package “evd” (STEPHENSON 2003). An example fit is shown in Figure 2a, which plots empirical and fitted distributions of S_{\max} for the cc strategy with 500 RILs. To determine genomewide significance thresholds we used the upper tail quantiles corresponding to p-values of $g(5\%)$, $g(1\%)$, $g(0.1\%)$ and $g(0.01\%)$, where $g(x) = 1 - (1 - x)^{1/20}$ Bonferroni corrects for the fact we simulate only one chromosome per animal but set thresholds as if there were twenty (ie, $g(5\%) \approx 0.26\%$).

Table 3 lists these significance thresholds calculated for all combinations of strategies and number of RILs for the case of a total background effect of $\theta_B = 45\%$.

We found that, for a given number of null simulations, estimating thresholds via a GEV model is more efficient than the traditional approach of taking empirical quantiles from the observed c.d.f of S_{\max} . To illustrate this point, in Figure 2b shows how threshold estimates from both methods vary with the number of null simulations available.

Assessing the efficacy of detection and fine-mapping

For a given breeding protocol and QTL system, we calculate two main quantities from 1,000 independent simulations: power to detect and location error. “Power to detect” is the probability that a QTL is detected at a given genomewide significance level α , defined as the proportion of trials in which the highest scoring locus exceeds the α -threshold. “Mean location error” is the mean absolute distance of the predicted QTL from the actual QTL in trials where the predicted QTL reaches α -level significance. We investigated measuring accuracy using root mean squared error but found its sensitivity to outliers (i.e. false positive QTL detection events) produced misleading results.

Results

Single marker association compared to HAPPY for detecting QTLs

We first investigated the effects of algorithm choice (SMA versus HAPPY) and SNP ascertainment. Two graphs in Figure 3 plot the performance of HAPPY and single marker association in fine-mapping a 5% QTL segregating in a population of 1,000 mice derived from 500 Collaborative Cross RIs (ie, with 2 replicate mice per RI strain). Note that here we

20

define a 5% QTL as one that accounts for 5% of the phenotypic variance among the eight founder strains, and that we include additional unobserved QTLs to make up the additive genetic variance to 50% (see Methods).

Figure 3a shows the power to detect a significant QTL. The orange and light green lines show the performance of SMA and HAPPY in the case where the QTL is an unaltered SNP drawn from the original GNF marker set. The dark red and dark green lines show what happens when the SNP chosen to be the QTL has had its strain distribution pattern (SDP) shuffled with respect to the founders in advance of the simulation and representing a more likely scenario (see Methods for details). In the first case, SMA does better than HAPPY, detecting the QTL with higher power at all thresholds. But if the SDP of the QTL is shuffled, then single marker association fails whereas HAPPY remains powerful. Similarly Figure 3b, which plots the location error against significance threshold, tells a similar story: when the QTL is an unaltered SNP, both HAPPY and single marker association predict its location with sub-centiMorgan accuracy. But when the alleles of the QTL have been shuffled, the location error of SMA predictions rises dramatically whereas those of HAPPY are almost unchanged. Hereafter, we consider only shuffled QTLs.

The graphs in Figure 4 plots the location error of HAPPY predictions against those of SMA in 1,000 simulations. For F2 populations (Figure 4a) the two methods are similar, with poor

localization (where the peak is more than 15 cM away from the true location) slightly more in HAPPY than in SMA (3.1% v 1.2% of cases). However, for CC populations (Figure 4b) HAPPY localises the QTL as accurately as before but SMA misses the QTL by >15 cM far more often (2.8% v 10.1% of cases). Hereafter, we consider only mapping predictions made by HAPPY for all strategies except BC and F2, for which we consider predictions made by both HAPPY and SMA.

Variation in haplotype block length

Table 4 gives summary statistics of distributions of the lengths of unrecombined haplotype blocks produced by the breeding strategies. Statistics were calculated from 1,000 simulations of 1,000 animals using the artificial fully informative marker set described in Methods. . The estimates have a slight upward bias because double recombinants returning to the same founder that occur between markers evade detection.. Despite this, the observed distributions were close to the exponential distribution predicted by theory. As expected, the block length shortens with more generations. For instance, within the group of CC strategies, cc, which has no selection or maintenance, has the longest blocks whereas cc08, which has two years of maintenance, has the shortest. Strategies rihs05 and rihs10 have block lengths comparable to cc04 and cc08, and this owes much to their similar maintenance periods. Although the maintenance periods of cc04 and cc08 are also close to those of riail08 and riail13, these RIAIL strategies have longer block lengths because, with two founder haplotypes rather than eight, double recombinants are always invisible and many recombinations are uninformative.

The shortest mean block length is seen in rihs30 at 4 cM. However, the block lengths of cc04 and rihs10 are not much longer, indicating that maintenance beyond a couple of years is subject to diminishing returns.

Single QTL: detection and resolution

We simulated a single additive QTL in RI, backcross and F2 intercross populations using a fixed population size of 1,000 animals. The allelic value of the QTL was constant across designs and accounted for 5% of the phenotypic variance among the eight founders. For the RI designs, we varied the number of recombinant inbred lines from which the 1,000 animals were drawn. For example, 100 RILs corresponds to 10 replicates from each of 100 lines. The results are plotted in Figure 5 and tabulated in Table 5.

The first graph in Figure 5 shows the power to detect an effect at the 5% genomewide significance level plotted against the number of RILs (with non-RI strategies at 0 RILs). Power is defined here as the proportion of 1,000 trials in which the highest scoring locus exceeds a given significance threshold. The cc strategy (red line) illustrates the trend among RI strategies: the greater the number of lines, the more power to detect. There is consistent variation between RI strategies too: at 500 RILs, two way strategies (ie those involving only two progenitor strains; blue) are more powerful than eight way ones, and strategies with more recombination (eg, cc08 or riail13) do worse than their less recombinant counterparts.

At 1,000 RILs, the eight way strategies catch up. Overall, BC and F2, two-strain strategies with minimal recombination, have the greatest power to detect QTL.

The second graph in Figure 5 shows the mean location error for detected QTLs. Whereas each point in the first graph was based on 1,000 simulations, in the second graph each point is based on only those simulations that were significant. For instance, the estimate for cc at 100 RILs (leftmost red point) is an average over only 40 simulations and so will have greater variance than the other plotted points. Nonetheless, the cc strategy (red) clearly shows how accuracy improves with increasing numbers of RILs. In particular, its elbow bend at 500 lines (1.1 cM) characterizes a process of diminishing returns where, past that point, modest gains hardly justify additional RILs. The remaining trends are the reverse of the first graph: more strains contributing to the cross and greater recombination give rise to sharper localization of the QTL, and this is most salient at 1,000 RILs, where the best performer, cc08, reaches 0.58 cM. F2 and BC deliver more accurate localization with SMA (open circles) than HAPPY (filled circles) but nonetheless lag behind the ≥ 500 RIL strategies.

Figure 5 does not show results from the RIHS strategies because their power was too low to allow meaningful estimates of location error. Echoing the trend above, power and degree of recombination were inversely related with 1,000 RILs of rihs05 detecting 14.1% of QTLs,

rihs10 detecting 2.2% and rihs30 detecting 0.5% (see Table 5). The explanation appears to be genetic drift.

Genetic drift of the foreground QTL

By definition the QTL segregates in the progenitor strains and has an allele frequency of 0.125, 0.25, 0.375 or 0.5, depending on how many founders carry its high allele. During breeding, the frequency drifts unpredictably, affecting the, QTL effect size and detectability in the final mapping population. Table 6 describes the genetic drift of the QTL under detection for the different breeding strategies and for different starting allele frequencies. We do not list the mean of the QTL allele frequency in the simulated mapping populations, since an average hides the underlying fluctuation. Rather, we give the standard deviation of allele frequencies, listing strategies prone to wide fluctuations first. The order of strategies closely matches the reversed order of power to detect a 5% QTL. Notably, rihs30 is the only breeding strategy where the allele frequency is so unstable as to drift to fixation occasionally.

QTL effect size

We investigated the ability of the cc strategy with 500 RILs to map QTLs of greater effect. Figure 6 shows detection and mapping of QTLs with effect sizes 5%, 10%, 20%, 30% and 40% using both HAPPY and SMA. In all cases, heritability was set at 50% with background QTLs making up the remaining genetic variance. Figure 6a shows that effect sizes of 10% or greater are uniformly detected with HAPPY (filled circles). With SMA (open circles),

guaranteed detection occurs for effect sizes of 30% or greater. The HAPPY line in Figure 6b is asymptotic, with the mean localization error at 0.24 cM for a 40% QTL (note: the average spacing between markers in the marker set is 0.19 cM). The localization by SMA shows small but consistent improvement with QTLs of effect size 10% or greater, yet never falls below the 2 cM level.

Epistasis

We next simulated a foreground QTL under strong epistatic control from an unlinked masking QTL lying on an ungenotyped chromosome (see Methods) and tried to detect and map that foreground QTL in a population of 1,000 animals. The combined effect size of the foreground and masking QTL was 10% in the founder population, with background QTLs making up the genetic variance up to 50%. The results are plotted in Figure 7 and tabulated in Table 7.

The first graph in Figure 7 shows the power to detect for the different breeding strategies. It exhibits the same trends as the detection graph in Figure 5 but with all points depressed by about 40%. The top performer is BC using SMA, which achieves greater than 90% power. The RI strategies lag far behind, with the best two way RI (riail08) reaching just above 60% and the best eight way strategies (hardly distinguishable cc and cctop10) coming close to 55% at 1,000 RILs. As before, the power of the RIHS strategies was too low to merit their

inclusion in this figure, with 1,000 RILs of rihs05, rihs10 and rihs30 achieving 9%, 1.9% and 0.5% respectively. The second graph in Figure 7 plots location error and tells a similar story to its Figure 5 counterpart, but with all strategies locating the QTL with roughly twice the error. RI strategies with ≥ 500 RILs outperform F2 and BC strategies, though in the case of BC mapped with SMA this is by a small margin. The elbow bend of the cc line (500 RILs, 1.7 cM) shows, as before, that increasing the number of lines beyond 500 delivers only modest gains in resolution. The most accurate strategy and the only one that reaches below the 1 cM mark is cc08 (0.98 cM).

Discussion

The collaborative cross is a highly attractive proposition because a large set of recombinant inbred strains are a powerful resource for investigating the genetic basis of complex traits, making it possible to map genetic effects into small regions and investigate gene by gene and gene by environment interactions. One drawback to the proposal is that the cross might have to be extremely large to meet some of its objectives: initial suggestions were that 1,000 strains would be required. Our simulation results for QTL mapping are more optimistic.

We explored the power and mapping resolution of CC strategies for identifying QTLs and compared the results to classical F2 intercross and BC approaches, as well as two other RI

methods: one using RIs derived directly from HS animals and the other from an advanced intercross. We found a number of striking trends that together characterize the relationship between power, accuracy and aspects of breeding design in two way and eight way crosses.

First, the increase in power and resolution that accompanies an increase in RI lines (Figure 5). We used a fixed population size of 1,000 animals regardless of how many RI lines were included. This meant we implicitly modeled the trade off between replication and genetic diversity, since one is bought at the cost of the other (see (BELKNAP 1998) for a theoretical treatment).

Second, there was tendency for more highly recombinant strategies to detect the QTL less readily but, once detected, map it more accurately. Higher recombination produces a finer grain haplotype mosaic that, by its reduced linkage disequilibrium between neighbouring markers, allows sharper localization. Most strategies achieve higher recombination through increased generation time (the exception being cctop10, which undergoes specific selection for recombination during breeding). More generations means greater opportunity for genetic drift and hence more cases of some QTL allele frequencies dropping to undetectable levels (FALCONER and MACKAY 1996). This explains the close resemblance between the order of strategies in Table 6 and their relative power in Figure 5 and Figure 7. Replication accentuates these undesirable effects by reducing the effective population size. Conversely,

increasing the number of lines stabilizes the allele frequencies and resulting in a convergence of data series as seen in the power graphs of Figure 5 and Figure 7.

Third, our simulations suggest greatest resolution is achieved through a hybrid collaborative cross strategy in which two maintenance years are inserted between mixing and inbreeding (cc08). In both simulations of single and epistatic QTLs, this strategy delivered sub-centiMorgan mapping with 1,000 RILs and only slightly less accurate predictions with 500 RILs. The standard cc localized with about 0.5 cM more error on average. However, the advantage of cc08 must be weighed against its lower power and the additional time and cost it would incur in stock creation. We acknowledge the precision of our QTL localizations was limited by the density of markers in the GNF set. Notably this limit caused the apparent asymptote in Figure 6 (see Results). We chose the GNF set because it was the best available at the time of simulation. Since then, however, there have been and will continue to be denser marker sets available. It is possible that performing the simulations with more closely-spaced markers may uncover a stronger advantage to using greater numbers of RILs than is suggested by Figure 5.

Our conclusions deserve certain qualifications, both general and specific in nature. On a general level, we chose a 5% QTL as a standard, but this may have provided a particularly stringent target, thereby making our results more conservative than necessary. Though a

recent review surveyed the effect sizes of reported QTLs phenotypes mapped using inbred crosses and found the mean was closer to 6% (FLINT *et al.* 2005), in fact a smaller effect size is likely to be more typical value. That is partly because most mapping studies over-estimate QTL effect sizes (due to their small sample size) and partly because experiments that dissect QTLs so often find that a single QTL is due to the combined effects of a number of physically linked QTLs of small effect (FLINT *et al.* 2005).

We should also point out that our results do not tell us about the ability of the strategies to detect epistasis. Rather, our concern was to map a QTL in the presence of epistasis. In our simulations the extent to which the epistatic QTL remains detectable depends on the minor allele frequency of the second (masking) locus, which in turn depends on how well the chosen breeding strategy preserves heterogeneity. We may therefore underestimate the power and resolution attainable in practice when all chromosomes can be analysed simultaneously.

A further general qualification is that our assessment of the trade off between replication and genetic diversity, which affects the advantage of adding RI lines, is crucially dependent on the level of background genetic variation, in the case of mapping a 5% QTL, we have set at 45%. Replication squeezes out environmental noise. Phenotyping an individual once with environmental error is always going to be less accurate than phenotyping ten identical individuals and taking their mean. That difference in accuracy is related to the size of the

environmental component. For instance, if variation in the trait were entirely genetic, replication could confer no advantage. During mapping we take into account whether the population contains replicates. If not, we regress on single animals. But if there are replicates, we regress on strain means. Doing so amounts to deriving a pseudo-population of fewer animals with a QTL of greater effect size. How does this influence our results?

Background genetic variation, QTL effect size and replication are related as follows. Let θ_Q be the effect size (ie, proportion of variance explained) in the mapping population of primary QTL, θ_B the summed effect size of all background QTLs and θ_E be the remaining environmental variance, where $\theta_E = 1 - (\theta_Q + \theta_B)$, and suppose the population of size N comprises N/n groups of n replicates. Then regressing on strain means is equivalent to mapping in a pseudo-population of N/n animals where the effect size of the primary QTL is now

$$\theta_q = \frac{\theta_Q}{\theta_Q + \theta_B + \frac{\theta_E}{n}}.$$

For instance, consider a 5% QTL with no background QTLs segregating in a population of 1000 genetically distinct animals (ie, $\theta_Q=0.05$, $\theta_B=0$, $N=1000$, $n=1$). Then consider the same QTL in a population that includes replicates: 1000 animals drawn in equal measures from 100 RILs (ie, $\theta_Q=0.05$, $\theta_B=0$, $N=1000$, $n=10$). Both instances describe a 5% QTL in a

population of 1000 individuals. But by averaging phenotypes across genetically identical animals, and thereby exploiting known population structure, the second instance is more usefully viewed as a 34% QTL in a pseudo-population of 100 “averaged” individuals.

How does inclusion of background QTLs change this? In our simulations background QTLs account for 45% of the phenotypic variance, such that the total heritability is 50% ($\theta_Q=0.05$, $\theta_B=0.45$). Without exploiting population structure, both instances still describe a 5% QTL segregating in 1000 individuals. However, treating the second instance as a pseudo-population of 100 now delivers an effect size of only 9%, showing a diminished advantage of replication in the face of a strong genetic background.

We chose a 50% heritable trait because it accorded with the estimated heritability of many physiological and behavioural traits. We found that power rose consistently with the number of lines used but that accuracy levelled off after 500 RILs, suggesting 500 may be an economic optimum at which acceptable power is accompanied by fine localization. However, we note that if the trait were less heritable, this optimum would be lower; if the trait were more heritable, it would be higher.

It is noteworthy that the BC and F2 apparently performed better than many of the RI strategies, particularly in the presence of epistasis and when the number of RI lines was low.

The main advantage of the RI is the much better mapping resolution achievable with ~500 lines. However, the two-way strategies F2, BC and RIAIL owed some of their success to strain ascertainment bias. Selecting only founder pairs in which the QTLs segregate conferred several advantages over the eight-way strategies. First, it ensured QTL allele frequency in the founding population was always 50% (while it could be as low as 12.5% in an eight way cross). Consequently, not only did the marker allele uniquely determine strain origin, but also the effect size of the QTL was often amplified. We set the allelic value (the amount that an allele adds to the phenotype), to be constant across designs and calibrated it so it would account for 5% of the variance in a population of the eight founders. Among the founders, most QTLs had a minor allele frequency of less than 50%. However, in the two-way strategies, the QTL frequency rises to 50%, which increases the variance attributable to it and thus its effect size. Moreover, because some background QTLs that segregate among eight founders will inevitably not segregate between two founders, the amount of background variation will be diminished. Note this is somewhat offset by the fact that the same will be true of non-QTL markers, fewer of which will be polymorphic, leading to coarser-grain mapping.

These are important consequences of using simpler breeding programmes. We could have set the effect size of the QTL according to the mapping population, but that would have required varying the allelic value across designs; an unfair depiction of biological reality. Nonetheless,

there is one advantage of eight way strategies our simulations mask: namely that a QTL is more likely to segregate among eight strains than two. We have assumed the QTL is known to segregate between the two chosen strains. Although pairs of strains are rarely chosen for a cross at random, it is a goal of the collaborative cross to make that decision less crucial to success of a mapping experiment. We discuss this point more fully in the Appendix and detail how it upwardly biases our assessment of power and accuracy for two way strategies.

The F2 and BC also have inherent disadvantages against RI strategies, two way or otherwise. Because we simulate dominance effects, homozygous individuals lie at the extremes of the phenotypic spectrum. That means the genetic variance of a QTL in an RI population will be greater than in a F2 population, which has heterozygotes, and particularly in a BC population, which lacks one homozygote class (BELKNAP 1998; FLINT *et al.* 2005). However, our inclusion of background QTLs moderates these effects.

The poor performance of the RIHS strategies is also thanks partly to genetic drift, but the explanation is slightly more complex and relates our inclusion of background QTLs. The simulated RIHS design was faithful to the construction of the existing Northport HS. In particular, the number of mating pairs in the mixing stage of breeding was set at 50. In the case of rihs30, that introduced a dangerous bottleneck that allowed fixation or near fixation at many loci. In simulations mapping a foreground QTL that affects a monogenic quantitative

trait we have found increasing the number of mating pairs to 150 recovers some power and accuracy (data not shown). But in the presence of background QTLs more extreme increases are necessary.

The rihs08 and rihs04 strategies echo these problems to a lesser extent. But the impact of the maintenance bottleneck in RIHS runs deeper. Table 3 lists 5% significance thresholds for different strategies. The thresholds for the rihs30 strategy are excessively high because the rihs30 population is quasi-inbred. Many markers on the genotyped chromosome (chromosome 1) are fixed or nearly fixed, and many of the background QTLs (on chromosomes 2-7; see Methods) have drifted far from their starting frequencies of 50% to have highly unequal effects. In our null simulations of rihs30, where the foreground QTL has zero effect, it is likely that some of the genotyped markers are in strong linkage disequilibrium (LD) with background QTLs of amplified effect. Because these markers act as surrogates for a true QTL, they will produce good regression fits even in the absence of a foreground QTL and, as a result, cause threshold estimates based on that null distribution to soar.

Moreover, increasing the number of lines from 500 RILs to 1,000 RILs will make matters worse. In the CC strategies, RILs are either drawn from a vast pool of genetically different individuals from an expansive mixing phase (as in cc and cctop10) or a large maintenance

population. In the RIHS strategies, RILs are always drawn from a small maintenance population. Therefore, increasing the number of RILs releases little extra genetic variation but does increase the apparent sample size and therefore the significance of the observed association. Note that RIHS performed far worse here than in simulations reported previously (VALDAR *et al.* 2003). The reasons are twofold. First, in those earlier simulations we did not include background QTLs and so avoided the pitfalls described above. Second, in those simulations we fixed the effect size of the QTL relative to the mapping population, which meant QTLs drifting to low frequency were given commensurately larger allelic values in order to produce a 5% effect. By contrast, our current simulations calibrate the allelic value in the founders and fix it thereafter such that the effect size of the QTL changes as its allele frequency drifts through breeding. Thus, our current simulations implicitly reward strategies that stem genetic drift and penalize those that encourage it.

In summary, we find that for QTL detection and high resolution mapping, the CC would not greatly benefit from increasing the number of inbred strains above 500. In the presence of epistatic QTLs, additional RIs certainly help, particularly with detection, but the gains are relatively modest. Given the cost and logistical difficulties inherent in creating, maintaining and distributing a very large set of RIs, it is hard to justify the routine use of a CC larger than 500 for QTL mapping, although there may be advantages using additional lines for other experimental purposes, such as expression analyses. The results presented here are not

specific to mice, and so will be of interest to geneticists designing panels of RIL for other model organisms and plants.

Finally, we found that no strategy described here will definitively identify a gene. The mapping resolutions are still insufficient to achieve that goal. In the best scenario we achieve using an eight-way strategy, the location error is just under 0.6 cM (obtained by two additional years of strain mixing in the CC and using 1,000 RIs). Note, this is not the same as a confidence interval. Rather, it means that if we were to map a QTL, the gene would lie on average 0.6 cM away from the highest association peak. Since the mean gene density in the mouse genome is about 10 genes per Mb, ideally we need a method with a resolution of 100 kb (equivalent to a location error of 0.05 cM).

Appendix

The success of two way strategies in our simulations is partly thanks to strain ascertainment bias. For those strategies, we selected founder pairs in which the QTLs segregate, since to do otherwise would have meant simulating no QTL for most cases. However, this means our assessment of power for these strategies is upward biased.

It is instructive to consider what would happen if, rather than ascertain strains, we choose the founder pair randomly from the eight. How would this affect power and resolution? The following reasoning assumes the QTL is one of the GNF SNPs and segregates among the eight founders. Power to detect describes a probability, $P(\text{detect})$. We can split this probability as follows:

$$P(\text{detect}) = P(\text{detect} | \text{segregates})P(\text{segregates}) + P(\text{detect} | \sim \text{segregates})P(\sim \text{segregates})$$

Where “segregates” means the QTL segregates and “ \sim segregates” means it does not. Clearly, $P(\sim \text{segregates}) = 1 - P(\text{segregates})$. Also the $P(\text{detect} | \sim \text{segregates})$, the probability to detect a QTL given it is fixed, is equivalent to the significance threshold:

$$P(\text{detect}) = P(\text{detect} | \text{segregates})P(\text{segregates}) + (0.05)(1 - P(\text{segregates}))$$

In the simulations performed here, we report only cases where the QTL segregates and so we report the power as $P(\text{detect} | \text{segregates})$. In the case of an eight way strategy, $P(\text{segregates})=1$, and so $P(\text{detect} | \text{segregates}) = P(\text{detect})$. However, for a two-way strategy without strain ascertainment, $P(\text{segregate})$, is less than 1 and so $P(\text{detect}) \neq P(\text{detect} | \text{segregate})$.

We can work out $P(\text{segregates})$ for a single SNP as follows. If the high allele exists in b out of 8 strains, then the probability of choosing two strains but only one high allele is

$$P(\text{segregates} | b) = \frac{2}{7}b - \frac{1}{28}b^2. \text{ Averaging this over all } m \text{ GNF SNPs gives}$$

$$\begin{aligned}
P(\text{segregates}) &= \sum_i^m P(\text{segregates} | b_i) P(i) \\
&= \frac{1}{m} \sum_i^m P(\text{segregates} | b_i) \\
&\approx 0.471
\end{aligned}$$

Where b_i is the value of b for marker i and $P(i)$ is the probability of choosing that marker.

Therefore, for the two-way crosses, the power to detect without strain ascertainment is

$$P(\text{detect}) = P(\text{detect} | \text{segregates})(0.471) + (0.05)(0.529)$$

which would mean an F2, BC or RIAIL whose power was 100% with strain ascertainment would only have about 50% power without this advantage. This reduction is even more pronounced in the epistatic case. There, further strain selection to ensure segregation of the unlinked QTL results in $P(\text{segregates})=0.269$, which reduces an apparent 100% detection to about 30%.

In summary, if we were to simulate the extreme opposite case of no strain ascertainment, for two-way strategies power would be about half that reported here when mapping a single QTL, and less than a third in the case of epistasis.

Acknowledgements

We thank Ken Paigen and Karl Broman for helpful discussions. This work was funded by a grant from The Wellcome Trust.

References

- BELKNAP, J. K., 1998 Effect of within-strain sample size on QTL detection and mapping using recombinant inbred mouse strains. *Behav Genet* **28**: 29-38.
- BOUCHER, W., and C. W. COTTERMAN, 1990 On the classification of regular systems of inbreeding. *J Math Biol* **28**: 293-305.
- CABALLERO, A., and M. A. TORO, 2000 Interrelations between effective population size and other pedigree tools for the management of conserved populations. *Genet Res* **75**: 331-343.
- CHURCHILL, G. A., D. C. AIREY, H. ALLAYEE, J. M. ANGEL, A. D. ATTIE *et al.*, 2004 The Collaborative Cross, a community resource for the genetic analysis of complex traits. *Nat Genet* **36**: 1133-1137.
- COLES, S., 2001 *An introduction to statistical modeling of extreme values*. Springer-Verlag, London.
- DARVASI, A., and M. SOLLER, 1995 Advanced intercross lines, an experimental population for fine genetic mapping. *Genetics* **141**: 1199-1207.
- DEMAREST, K., J. KOYNER, J. MCCAUGHRAN, JR., L. CIPP and R. HITZEMANN, 2001 Further characterization and high-resolution mapping of quantitative trait loci for ethanol-induced locomotor activity. *Behav Genet* **31**: 79-91.
- DUDBRIDGE, F., and B. P. KOELEMAN, 2004 Efficient computation of significance levels for multiple associations in large studies of correlated data, including genomewide association studies. *Am J Hum Genet* **75**: 424-435.
- FALCONER, D., and T. F. MACKAY, 1996 *Introduction to quantitative genetics*. Longman Group Ltd.
- FLINT, J., W. VALDAR and R. MOTT, 2005 Experimental strategies for mapping and cloning quantitative trait genes in rodents. *Nat Rev Genet* **6**: 271-286.
- KIMURA, M., and J. CROW, 1963 On the maximum avoidance of inbreeding. *Genetical Research* **4**: 399-415.
- LYNCH, M., and B. WALSH, 1998 *Genetics and analysis of quantitative traits*. Sinauer Associates, Inc, Sunderland, Massachusetts, USA.
- MCCLEARN, G. E., J. R. WILSON and W. MEREDITH, 1970 The use of isogenic and heterogenic mouse stocks in behavioral research, pp. 3-32 in *Contribution to behavior*

- genetic analysis. The mouse as a prototype*, edited by L. G and T. D. D. Appleton Century Crofts, New York.
- MOTT, R., C. J. TALBOT, M. G. TURRI, A. C. COLLINS and J. FLINT, 2000 A method for fine mapping quantitative trait loci in outbred animal stocks. *Proc Natl Acad Sci U S A* **97**: 12649-12654.
- PLETCHER, M. T., P. MCCLURG, S. BATALOV, A. I. SU, S. W. BARNES *et al.*, Use of a dense single nucleotide polymorphism map for in silico mapping in the mouse. *PLoS Biology* **2**.
- STEPHENSON, A., 2003 evd: Extreme Value Distributions. *R News* **2**: 31-32.
- TALBOT, C. J., R. A. RADCLIFFE, J. FULLERTON, R. HITZEMANN, J. M. WEHNER *et al.*, 2003 Fine scale mapping of a genetic locus for conditioned fear. *Mamm Genome* **14**: 223-230.
- THE COMPLEX TRAIT CONSORTIUM, 2003 A Collaborative Cross for High-Precision Complex Trait Analysis v31, pp. in <http://www.complextait.org/>.
- VALDAR, W. S., J. FLINT and R. MOTT, 2003 QTL fine-mapping with recombinant-inbred heterogeneous stocks and in vitro heterogeneous stocks. *Mamm Genome* **14**: 830-838.
- WANG, X., I. LE ROY, E. NICODEME, R. LI, R. WAGNER *et al.*, 2003 Using advanced intercross lines for high-resolution mapping of HDL cholesterol quantitative trait loci. *Genome Res* **13**: 1654-1664.

TABLES

Type of strategy	Strategy Name	Mixing	Number of maintenance generations	Inbreeding for 20 generations	Total number of generations
CC	cc	disjoint diallel	0	yes	3+0+20=23
CC	cctop10	disjoint diallel*	0	yes	3+0+20=23
CC	cc04	disjoint diallel	4	yes	3+4+20=27
CC	cc08	disjoint diallel	8	yes	3+8+20=31
RIHS	rihs05	simple	5	yes	3+5+20=28
RIHS	rihs10	simple	10	yes	3+10+20=33
RIHS	rihs30	simple	30	yes	3+30+20=53
RIAIL	riail08	simple	7	yes	1+7+20=28
RIAIL	riail13	simple	10	yes	1+10+20=33
BC	bc	simple	1**	no	1+1+0=2
F2	f2	simple	1	no	1+1+0=2

Table 1. Strategies simulated.

Key: Collaborative Cross (CC); recombinant inbred heterogeneous stocks (RIHS); recombinant inbred advanced intercross lines (RIAIL); F2 intercross (F2). See Methods for descriptions of "disjoint diallel" and "simple" mixing.

*** Strategy includes selection: only the most highly recombinant 10% of animals from the four-way (F2) cross participate in the eight-way (F3) cross.**

****Maintenance involves mating with one of the parental lines in the case of BC**

QTL system	Genotype	Genetic effect (arbitrary units)
Single	aa	0
	aA	5
	AA	10
Epistatic	aabb	0
	aAbb	0
	AAbb	0
	aaBb	0
	aABb	5
	AABb	10
	aaBB	0
	aABB	10
	AABB	20

Table 2. Mean effects for each multilocus genotype in simulated additive and epistatic QTL systems. In non-epistatic simulations, the genetic effects of background and foreground QTLs follow the additive scheme. In epistatic simulations, background QTLs act additively but the foreground QTL and the unlinked masking QTL act according to the epistatic scheme. Note that, although listed, heterozygous genotypes are extremely rare in inbred RIAIL, RIHS and CC populations. For this reason, we excluded dominance effects from either model.

Strategy	Estimated 5% genomewide significance threshold (observed quantile in 1000 null simulations)		
	0 RILs	500 RILs	1000 RILs
cc*	-	4.51 (4.69)	4.49 (4.19)
cc04	-	4.75 (5.22)	4.59 (4.41)
cc08	-	4.88 (4.36)	4.95 (4.58)
cctop10	-	4.65 (4.81)	4.4 (5.05)
riail08	-	6.2 (6.46)	6.96 (7.02)
riail13	-	7.39 (8.55)	8.83 (9.87)
rihs05	-	9.15 (9.52)	13.93 (14.15)
rihs10	-	14.33 (12.61)	22.75 (19.48)
rihs30	-	28.22 (19.95)	41.49 (29.77)
bc	5.03 (4.57)	-	-
f2	5.2 (5.08)	-	-

Table 3. Significance thresholds for HAPPY scores (ie, $-\log_{10}$ F-test P values) calculated for different combinations of breeding strategy and number of RILs. Each threshold is calculated from a Generalized Extreme Value (GEV) distribution that has been fit to the maximum scores obtained from 1,000 simulations of mapping a zero-effect QTL (ie, 1,000 null simulations). Null simulations listed here had the level of background genetic variation at 45% and so apply only to mapping of a 5% foreground QTL. In brackets are thresholds calculated directly from the empirical cumulative distribution of null maxima (shown for comparison). Note that correction for anticipated multiple chromosomes means the 5% genomewide level corresponds to a chromosomewide significance level of ~0.26%.

* Additional simulations were performed for cc, with results (written as number of RILs = power (location error)): 100 RILs=4.62 (4.49), 200 RILs=4.62 (5.25), 250 RILs=4.49 (4.35), 333 RILs=4.43 (4.55).

Strategy name	Apparent unrecombined blocklength in cM	
	Mean	S.D.
cc	11.51	12.18
cctop10	10.09	10.46
cc04	8.56	8.89
cc08	6.91	7.12
rihs05	8.66	9.05
rihs10	6.85	7.11
rihs30	4.00	4.15
riail08	12.90	13.66
riail13	10.47	11.14
BC	68.41	36.88
F2	36.84	26.79

Table 4. Unrecombined block length. Mean, Standard Deviation (S.D.) for each strategy are estimated from 1000 simulated populations of 1000 individuals. The standard error for the blocklength means are of the order 0.01.

Strategy	Power to detect QTL			Mean location error (cM)		
	0 RILs	500 RILs	1000 RILs	0 RILs	500 RILs	1000 RILs
cc*	-	0.67	0.94	-	1.14	0.87
cc04	-	0.63	0.91	-	0.92	0.71
cc08	-	0.49	0.82	-	0.96	0.58
cctop10	-	0.64	0.93	-	1.28	0.79
riail08	-	0.87	0.93	-	1.08	1.01
riail13	-	0.75	0.84	-	1.12	1.06
rihs05	-	0.19	0.14	-	2.20	0.89
rihs10	-	0.04	0.02	-	3.39	8.50
rihs30	-	0.00	0.01	-	0.85	1.18
bc	1.00	-	-	2.78	-	-
bc (SMA)	1.00	-	-	1.49	-	-
f2	0.96	-	-	3.64	-	-
f2 (SMA)	0.99	-	-	2.22	-	-

Table 5. Detection and mapping of a single 5% foreground QTL in 1,000 animals. See Figure 5 legend and Results for details.

***Additional simulations were performed for strategy cc with results (written as number of RILs=power (location error)): 100 RILs=0.04 (4.27), 200 RILs=0.14 (2.24), 250 RILs=0.28 (1.79), 333 RILs=0.47 (1.66).**

Strategy	Standard deviation of QTL minor allele frequency for starting frequency q				
	q=12.5%	q=25%	q=37.5%	q=50%	% Fixed
rihs30	9.5	11.0	10.2	8.7	0.9
rihs10	6.9	8.8	8.1	6.7	0
rihs05	6.2	7.4	7.7	5.5	0
riail13				5.3	0
riail08				4.5	0
ccall2yr	1.5	2.1	2.3	1.4	0
ccall1yr	1.4	1.9	2.1	1.3	0
cctop10				1.2	0
ccall				1.2	0
bc				0.9	0
f2				0.7	0

Table 6: Genetic drift of the foreground QTL in different 500 RIL breeding strategies (with f2 and bc included for comparison). Before breeding, the QTL has a starting minor allele frequency of q in the founders. During breeding that frequency fluctuates and in the mapping population it may be elevated or depressed. This table shows the extent of drift from founder to mapped individual, expressed as the standard deviation of allele frequency in the mapping population. The more drift, the greater the tendency for alleles to approach fixation.

Strategy	Power to detect QTL			Mean location error (cM)		
	0 RILs	500 RILs	1000 RILs	0 RILs	500 RILs	1000 RILs
cc*	-	0.29	0.54	-	1.72	1.34
cc04	-	0.25	0.47	-	1.34	1.07
cc08	-	0.18	0.40	-	0.98	0.82
cctop10	-	0.28	0.55	-	1.92	1.27
riail08	-	0.49	0.61	-	1.98	1.49
riail13	-	0.37	0.46	-	2.10	1.89
rihs05	-	0.11	0.09	-	3.01	2.25
rihs10	-	0.06	0.02	-	6.82	9.71
rihs30	-	0.00	0.01	-	-	2.13
bc	0.84	-	-	4.99	-	-
bc (SMA)	0.92	-	-	2.36	-	-
f2	0.60	-	-	4.78	-	-
f2 (SMA)	0.78	-	-	3.05	-	-

Table 7. Detection and mapping of an epistatic QTL in 1,000 animals. Tabulation of data in Figure 7. See figure legend and Results for details.

*** Additional simulations were performed for cc, with results (written as number of RILs = power (location error)): 100 RILs=0.02 (8.16), 200 RILs=0.05 (3.20), 250 RILs=0.12 (3.07), 333 RILs=0.19 (2.80)**




Figure 1 Components of a generalized breeding strategy. The first stage, mixing, combines different color-coded inbred strains to produce mice containing some genetic material from each founder. In the optional maintenance stage, mice are intercrossed to increase the number of recombination breakpoints. A subsequent inbreeding produces recombinant inbred lines.




Figure 2: Fitting a Generalized Extreme Value distribution (GEV) to the null distribution of maximum $-\log P$ for the cc strategy with 500 RILs. Graph (a) shows the empirical and fitted densities for $-\log P$ of the best fitting locus in each of 1000 simulations of a zero-effect QTL. Underneath the curves, a rug plot shows individual data points. Graph (b) plots chromosomewide and genomewide thresholds estimated by two methods: “Quantile” takes a suitable quantile of the empirical distribution (eg, the 5% chromosomewide threshold would equals the 95% quantile of the distribution of observed scores), whereas “GEV” estimates thresholds analytically via a fitted GEV. Estimates are plotted for 1,000 and 10,000 null simulations, with 10,000 expected to be the more accurate. The plot shows that whereas quantile estimates vary substantially with the number of simulations and especially for small significance levels, GEV estimates are relatively consistent.






Figure 3: The effects of marker ascertainment bias on mapping power and resolution. A diallelic 5% QTL was mapped in a population of 1,000 mice derived from 500 cc RIs, either using the original strain distribution pattern (green) or a shuffled SDP (red) for two mapping algorithm SMA and HAPPY. Shuffling removes any marker ascertainment bias. (a) Power to detect a significant QTL. (b) The location error for a predicted QTL.



(a)



(b)

Figure 4: Location error of predicted QTLs using HAPPY v single marker association (SMA) in 1,000 simulated mapping populations (a) 1,000 F2 (b) 1,000 individuals from 500 collaborative cross recombinant inbred lines






Figure 5 Detection and mapping a single additive 5% QTL using HAPPY (filled circles) and SMA (open circles, F2 and BC only) for different breeding strategies. Each data point is based on 1,000 simulations of mapping in 1,000 individuals. The horizontal axes indicate the number of RI lines in the cross. Zero represents backcross (BC) and F2 strategies; 100 (middle tick) refers to RI strategies with 10 replicates for each of 100 RI lines; 1,000 means one mouse for each of 1,000 RI lines. Note that a 5% QTL in an RI translates to a 2.56% QTL in an F2 and a 1.3% QTL in a BC (see text). The left graph shows power, ie, the probability that the highest scoring putative QTL achieves 5% genomewide significance. The right graph plots the location error, ie, the mean absolute discrepancy between the actual and predicted location of the QTL, for mapping experiments that achieved 5% significance or better.



(a)

(b)

Figure 6. Effect of varying QTL effect size on mapping strategy cc with 500 RILs. Results are shown for detection (a) and mapping (b) of a single additive QTL using HAPPY in a population of 1,000 mice derived from 500 Collaborative Cross (strategy cc) RI lines. Values are plotted for effect sizes 5%, 10%, 20%...90%.




Figure 7 Detection and mapping of an epistatic QTL. Graphs show results of finding a single QTL embedded within an epistatic system in which the second QTL is invisible, unlinked and has a masking effect, and where the broad sense heritability of the two QTLs is 10% (adjusted to 5.3% in F2 and 2.8% in BC to maintain constant allelic value). Results shown for HAPPY (filled circles) and SMA (open circles, F2 and BC only) for different breeding strategies. Keeping the population size at 1,000 but increasing the number of RI lines leads to increased power and accuracy. We illustrate this more fully for the cc strategy, reporting results for 100, 200, 250, 333, 500 and 1,000 RI lines.